

1.0 The Use of Enteral Nutrition vs. Parenteral Nutrition

Question: Does enteral nutrition compared to parenteral nutrition result in better outcomes in the critically ill adult patient?

Summary of evidence: There were 20 level 2 studies and one level 1 study (Woodcock et al) that were reviewed and meta-analyzed. In the Woodcock study, data from ICU patients only were abstracted and there were 11/38 patients that crossed over between EN and PN group after randomization. There have been two more recent, large RCTs, Harvey 2014 and Reignier 2017, which enrolled 2400 and 2410 patients, respectively, across 33 and 44 sites. Other more recent smaller trials included patients fasting for at least 14 days (Xi 2014), patients with moderate traumatic brain injury (Meirelles 2011) and patients with severe acute pancreatitis (Wang 2013, Sun 2013). A priori, we considered that the harmful effect of PN may be associated with relative overfeeding and hyperglycemia. Accordingly, we conducted a subgroup analysis to determine the effect of excess calories (PN compared to EN) and higher glucose levels (across groups). The Moore 1992 study, which had been included in the 2009 summary, was reviewed again and excluded since it reports results of a meta-analysis and the individual studies have been included. Given concerns about population in the Meirelles 2011 and Wang 2013 studies not being critically ill as no mention of ventilation status and some missing data in the latter study, a sensitivity analysis was also done excluding these two studies.

Mortality: In the two largest studies (Harvey and Reignier), there were no significant differences between the PN group and the EN group in 30 or 28 day mortality ($P = 0.57$ and 0.33 , respectively) or 90 day mortality ($P = 0.4$ and 0.28 , respectively) or hospital mortality ($P = 0.44$ and 0.25 , respectively). However, both studies showed a trend in the reduction in ICU mortality, favoring the PN group ($P = 0.13$ and 0.17 , respectively). When these data were aggregated with the other 17 studies reporting on mortality, there was no difference in overall mortality between the groups receiving EN or PN (RR 1.04, 95% CI 0.97, 1.12, $p=0.29$, test for heterogeneity $I^2=2\%$, figure 1). When the trials in which the PN group were fed more calories than the EN group were aggregated, there was no effect seen (RR 1.19, 95% CI 0.86, 1.64, $p = 0.30$, heterogeneity $I^2=31\%$; figure 1). Similarly, when the trials in which the PN and EN groups were fed isocalorically were aggregated, there was no effect on mortality (RR 1.02, 95% CI 0.92, 1.13, $p = 0.65$, test for heterogeneity $I^2=0\%$; figure 1). There was no difference in these subgroups ($p=0.39$; figure 1). In subgroup analysis comparing studies in which the PN group had higher blood sugars than the EN group to studies in which there was no difference in blood sugars, showed that increased mortality in the PN groups could not be explained by hyperglycemia (RR 0.93, 95% CI 0.30, 2.90, $p=0.90$, heterogeneity $I^2=0\%$; figure 2). In a sensitivity analysis excluding Meirelles 2011, Wang 2013, there was still no difference in mortality between groups (RR 1.04, 95% CI 0.97, 1.12, $p=0.21$, test for heterogeneity $I^2=0\%$, figure not shown). When data from the 6 studies reporting on ICU mortality were aggregated, there was no effect seen (RR 1.04, 95% CI 0.97, 1.12, $p = 0.28$, heterogeneity $I^2=0\%$, figure 3). There was also no effect seen when looking at subgroups where the PN group was fed more than the EN group and where the two groups were fed isocalorically ($p = 0.38$ and 0.71 , respectively, figure 3).

Infections: When the 12 studies which reported on patients with infectious complications were statistically aggregated, the meta-analysis showed that EN compared to PN was associated with a significant reduction in the incidence of infectious complications (RR 0.74, 95% CI 0.59, 0.91, $p=0.005$, heterogeneity $I^2=42\%$; figure 4). When the trials in which the PN group were fed more calories than the EN group were aggregated, EN compared to PN was also associated with a significant reduction in the incidence of infectious complications (RR 0.58, 95% CI 0.39, 0.88, $p=0.009$, heterogeneity $I^2=53\%$; figure 4). When the trials in which the PN and EN groups were fed isocalorically were aggregated, EN compared to PN had no effect on infectious complications (RR 0.94, 95% CI 0.80, 1.10, $p=0.44$, heterogeneity $I^2=0\%$; figure 4). There was a significant difference in these subgroups ($p=0.03$; figure 4). Another subgroup analysis showed that there was a trend between the increase in infections and hyperglycemia (RR 0.79, 95% CI 0.56, 1.11, $p=0.17$, heterogeneity $I^2=0\%$; figure 5). In a sensitivity analysis excluding Mereilles 2011, EN compared to PN was associated with a significant reduction in infectious complications (RR 0.66, 95% CI 0.50, 0.86, $p=0.003$, heterogeneity $I^2=38\%$, figure not shown).

LOS, Ventilator days: A total of 10 studies reported on hospital length of stay (in mean and standard deviation) and when the data were aggregated, enteral nutrition was associated with a trend towards a reduction in hospital LOS (WMD -3.12, 95% CI -7.43, 1.19, $p=0.16$, test for heterogeneity $I^2=96\%$; figure 6). Only 6 studies reported on ICU LOS (in mean and standard deviation) and when the data were aggregated, the use of EN was associated with a reduction in ICU LOS (WMD -2.12, 95% CI -4.20, -0.04, $p=0.05$, heterogeneity $I^2=94\%$; figure 7). A total of 5 studies reported on length of mechanical ventilation (in mean and standard deviation) and when the data were aggregated, there was a trend towards a reduction in ventilator days in the EN fed group (WMD -1.23, 95% CI -2.80, 0.34, $p=0.13$, heterogeneity $I^2=87\%$, figure 8).

Nutritional complications: Of the 13 studies that reported on nutritional intake, 5 found that PN was associated with a higher calorie intake (Rapp, Young, Moore, Kudsk, Woodcock (Blood sugar values in the Woodcock pertain to the entire group, not the ICU population), the remaining 8 reported no significant difference in intakes between the groups (Adams, Hadley, Cerra, Dunham, Borzotta, Kalfarentzos, Wang, Harvey). A total of 7 studies reported on hyperglycemia and in 4 of these, EN was associated with lower incidences of hyperglycemia compared to PN (Adams $p<0.001$), (Borzotta $p<0.05$, Kalfarentzos) (Mereilles $p<0.01$). Three studies showed no difference in blood sugars between the groups receiving EN and PN (Moore 1989, Rapp, Harvey). Four studies showed that EN was associated with an increase in diarrhea (Cerra $p<0.05$, Young, Kudsk $p<0.01$, Harvey) while one showed an association with EN and a reduction in diarrhea (Borzotta $p<0.05$) and one study showed no difference (Adam). One study showed reported higher increases in total protein and albumin levels in the EN group vs. PN group level ($p=0.032$ and $p=0.028$ respectively, Chen 2019)

Other Complications: EN was also associated with an increase in vomiting (Cerra $p < 0.05$), Harvey 2014 $p < 0.001$). One study found less favourable neurological outcome at 3 months ($p = 0.05$) in brain injured patients (Young, $p = 0.05$), though this significance disappeared after 6 months and 1 year. More overall nutrition related complications were noted in EN vs PN (Dunham). Seven studies reported on diarrhea. There were significant reductions in the incidence of hypoglycemia (44 patients [3.7%] vs. 74 patients [6.2%]; $P = 0.006$) in the parenteral group in the largest study (Harvey 2014)

Cost: Four studies reported a cost savings with the use of EN vs PN (Adams, Cerra, Borzotta and Kalfarentzos).

Quality of Life (QOL) Outcomes: In a second publication (Harvey 2016), quality of life from the Harvey 2014 study was reported. In the trial, the EuroQol 5-dimension (5-level version) questionnaire (EQ-5D-5L) and a Health Services Questionnaire (to evaluate health and nutrition related quality of life (QOL)) were completed at 90 days post randomization and 1 year post-randomization with survivors. At 90 days and 1 year post randomization, Harvey et al found that health components from the EQ-5D-5L questionnaire were similar between groups. The results for nutrition related QOL were reported on a scale from 1 (worst possible satisfaction) to 7 (best possible satisfaction). At 90 days post-randomization, there was no difference in the mean response between the PN (mean (SD) of 5.2 (1.6, $n=405$)) and EN groups (5.1 (1.7, $n=378$)) (mean difference 0.10, 95% CI, -0.14, 0.33, $p=0.43$) (data not shown in table). At 1 year, there was also no significant difference (5.3 (1.6) in the PN group ($n=338$) vs 5.4 (1.6) in the EN group ($n=322$), mean difference -0.10, 95% CI, -0.35, 0.14, $p=0.41$) (data not shown in table).

Conclusions:

- 1) The use of EN compared to PN has no effect on mortality in critically ill patients.
- 2) The use of EN compared to PN is associated with a reduction in the number of infectious complications in the critically ill in trials where patients in the PN group received more calories than in the EN group.
- 3) The use of EN compared to PN may be associated with a reduction in ICU LOS and ventilator days and hospital LOS. Significant heterogeneity limits the inferences from these aggregated analyses.
- 4) The use of EN compared to PN may not be associated with an improvement in calories due to underfeeding in both groups
- 5) The use of EN may be associated with increased episodes of vomiting.
- 6) There is no difference between EN and PN in terms of patient reported outcomes

Level 1 study: if all of the following are fulfilled: concealed randomization, blinded outcome adjudication and an intention to treat analysis

Level 2 study: If any one of the above characteristics are unfulfilled.

Table 1. Randomized studies evaluating EN vs PN in critically ill patients

Study	Population	Methods (score)	Intervention	Mortality # (%)†		Infections # (%)‡	
				EN	PN	EN	PN
1. Rapp 1983	Head Injured patients N=38 (<Ideal weight) Single-centre	C.Random: not sure ITT: no Blinding: no (4)	EN vs PN	9/18 (50)	3/20 (15)	NR	NR
2. Adams 1986	Trauma patients undergoing laporotomy N=46 36/46 ICU patients Single-centre	C.Random: not sure ITT: yes Blinding: no (8)	EN vs PN	1/23 (4)	3/23 (13)	15/23 (65)	17/23 (74)
3. Young 1987	Brain injured patients N=58 (N=51 randomized) Single-centre	C.Random: not sure ITT: no Blinding: no (6)	EN vs PN	10/28 (36)	10/23 (43)	5/28 (18)	4/23 (17)
4. Peterson 1988	Critically ill patients with abdominal trauma N=59 Single-centre	C.Random: not sure ITT: no Blinding: no (5)	EN vs PN	NR	NR	2/21 (10)	8/25 (32)
5. Cerra 1988	ICU patients post sepsis N=70 (hypermetabolic patients) Single-centre	C.Random: not sure ITT: no Blinding: no (2)	EN vs PN	ICU 7/31 (22)	ICU 8/35 (23)	NR	NR
6. Moore 1989	Abdominal trauma patients N=75 Single-centre	C.Random: yes ITT: no Blinding: no (10)	EN vs PN	NR	NR	5/29 (17)	11/30 (37)

7. Kudsk 1992	Abdominal trauma N=98 Single-centre	C.Random: not sure ITT: no Blinding: single (10)	EN vs PN	ICU 1/51 (2)	ICU 1/45 (2)	9/51 (16)	18/45 (40)
8. Dunham 1994	Blunt trauma N=38 Single-centre	C.Random: not sure ITT: no Blinding: no (8)	EN vs PN	1/12 (7)	1/15 (8)	NR	NR
9. Borzotta 1994	Closed head injury N=59 Single-centre	C.Random: not sure ITT: no Blinding: no (6)	EN vs PN	5/28 (18)	1/21 (5)	51/28 per group	39/21 per group
10. Hadfield 1995	ICU patients, mainly cardiac bypass N=24 Single-centre	C.Random: not sure ITT: no Blinding: no (7)	EN vs PN	ICU 2/13 (15)	ICU 6/11 (55)	NR	NR
11. Kalfarentzos 1997	Severe acute pancreatitis N=38 Single-centre	C.Random: not sure ITT: no Blinding: single (9)	EN vs PN	ICU 1/18 (6)	ICU 2/20 (10)	5/18 (28)	10/20 (50)
12. Woodcock 2001	Patients needing nutrition support N=562 ICU patients N=38 (all degrees of malnutrition) Single-centre	C.Random: yes ITT: yes Blinding: single (12)	EN vs PN	9/17 (53)	5/21 (24)	6/16 (38)	11/21 (52)

13. Casas 2007	Severe acute pancreatitis; ICU≥72 hrs N=22 Single-centre	C.Random: no/unsure ITT: Yes Blinding: No (8)	EN vs PN	Hospital 0/11 (0)	Hospital 2/11 (18)	1/11 (9)	3/11 (27)
14. Chen 2011	Elderly Patients in respiratory intensive care unit N=147 Single-centre	C.Random: Yes ITT: Yes Blinding: No (7)	EN vs PN	20-day 11/49 (22)	20-day 10/49 (20)	5/49 (10)	18/49 (37)
15. Meirelles 2011	Adult patients with moderate traumatic brain injury N=22 Single-centre	C.Random: No ITT: No Blinding: No (5)	EN vs PN	Unspecified 1/12 (8.3)	Unspecified 1/10 (10)	Total infectious complications 2/12 (16.7) Pneumonia (cases) 2/12 (16.7) Sepsis (cases) 0	Total infectious complications 4/10 (40) Pneumonia (cases) 2/10 (20) Sepsis (cases) 2/10 (20)
16. Wang 2013	Patients 18-45 years with severe acute pancreatitis N=183 Single-centre	C.Random: No ITT: No Blinding: Double (7)	EN vs PN	Hospital 3/61 (5)	Hospital 7/60 (12)	Pancreatic sepsis 13/61 (21) MODS 15/61 (24.6)	Pancreatic sepsis 24/60 (40) MODS 22/60 (36.7)
17. Sun 2013	Severe acute pancreatitis admitted to surgical ICU N=60 Single-centre	C.Random: No ITT: No Blinding: No (6)	EN vs PN	Hospital 2/30 (7)	Hospital 1/30 (3)	Pancreatic 3/30 (10) MODS 5/30 (17) SIRS 12/30 (40)	Pancreatic 10/30 (33) MODS 13/30 (43) SIRS 22/30 (73)

18. Harvey 2014	Adult patients admitted to a general ICU N=2400 Multi-centre	C.Random: Yes ITT: Yes Blinding: No (8)	EN vs PN	ICU 352/1197 (29.4) Hospital 450/1186 (37.9) 30-day 409/1195 (34.2) 90-day 464/1188 (39.1)	ICU 317/1190 (26.6) Hospital 431/1185 (36.4) 30-day 393/1188 (33.1) 90-day 442/1184 (37.3)	Total infectious complications 194/1197 (16.2)** Infectious complications per pt 0.21 +/- 0.5 Pneumonia 143/1197 (11.9) Bloodstream inf 21/1197 (1.8) Surgical inf 12/1197 (1.0)	Total infectious complications 194/1191 (16.3)** Infectious complications per pt 0.22 +/- 0.6 Pneumonia 135/1191 (11.3) Bloodstream inf 27/1191 (2.9) Surgical inf 10/1191 (0.8)
19. Xi 2014	ICU pts fasting at least 14 days, eligible for EN. Single Centre. N=45	C.Random: No ITT: Yes Blinding: No (7)	EN vs PN	28-day 0/22	28-day 0/23	Positive blood cultures 4/22 Sepsis 4/22 (17)	Positive blood cultures 0/23 Sepsis 5/23 (23)
20. Reignier 2017	Mechanically ventilated ICU pts receiving vasopressor support for shock. Multi-centre. N=2410	C.Random: Yes ITT: Yes Blinding: No (11)	EN vs PN	ICU 429/1202 (33) Hospital 498/1202 (36) 28-day 443/1202 (37) 90-day 530/1185 (45)	ICU 405/1208 (31) Hospital 479/1208 (34) 28-day 422/1208 (35) 90-day 507/1192 (43)	ICU acquired 173/1202 (14)	ICU acquired 194/1208 (16)
21. Chen 2019	Severe acute pancreatitis within 48 hrs admission Single centre N=140	C.Random: no ITT: yes Blinding: No (7)	EN vs PN EN via jejunostomy	Not specified † 10/70 (14.3%)	Not specified † 12/70 (17.1%)	NR	NR

C.Random: concealed randomization

* median/mean values, no standard deviation hence not included in meta-analysis

‡ refers to the # of patients with infections unless specified

** data on ICU patients/infections obtained directly from author

ITT: intent to treat

NR: not reported

† presumed hospital mortality unless otherwise specified

± () : mean ± Standard deviation (number)

reported data pertaining to ICU patients only

NS = not statistically significant

Table 1. Randomized studies evaluating EN vs. PN in critically ill patients (continued)

Study	LOS days		Ventilator days		Cost		Other	
	EN	PN	EN	PN	EN	PN	EN	PN
1. Rapp 1983	Hospital 49.4*	Hospital 52.6*	10.3*	10.4*	NR	NR	Mean Calorie Intake (kcal) in study 685 1750 p=0.001 Nitrogen Intake (gms) 4.0 10.2 p=0.002 N balance/24hrs, p=0.002 -17.6 -10.9 Hyperglycemia no difference between groups	
2) Adams 1986	ICU 13 ± 11 (19) Hospital 30 ± 21 (19)	ICU 10 ± 10 (17) Hospital 31 ± 29 (17)	12 ± 11 (17)	10 ± 10 (13)	\$1346/day	\$3729/day	Calorie Intake (kcal) in study 2088 2572 p=NS Caloric adequacy 73% 89% N balance/24 hrs -8.7+6.8 -4.1+4.6 Hyperglycemia (pt days) 24/242 (10) 49/220 (22) p<0.001 Line Problems 13/9 9/7 Diarrhea (days/pt) 3.5 3.8	
3. Young 1987	NR	NR	NR	NR	NR	NR	Calories ÷ BEE x 1.75 59% ± 5.13% 75.6% ± 4.26% p=0.02 Protein Intake (gm/kg/day) 0.91 ± 0.09 1.35 ± 0.12 p=0.004 Favourable Neurological Outcome (3 months) 17.9 % 43.5 % Diarrhea 23/28 (82) 13/23 (57)	

<p>4. Peterson 1988</p>	<p>ICU 3.7 ± 0.8 (21) Hospital 13.2 ± 1.6 (21)</p>	<p>ICU 4.6 ± 1.0 (25) Hospital 14.6 ± 1.9 (24)</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>	<p>Day 5 Calorie Intake (kcal) 2204 ± 173 2548 ± 85 P=0.04 Day 5 Nitrogen Intake (gms) 12.6 ± 1.0 14.8 ± 0.6</p>
<p>5. Cerra 1988</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>	<p>\$228 ± 59 /day</p>	<p>\$330 ± 61 /day</p>	<p>NON PROTEIN Calorie Intake 1684 ± 573 2000 ± 20 p=NS Protein g/d 80±26 88±20 N-balance/d -3.4±10 0.4±3.8 MOSF 7/31 (23) 7/35 (20) Diarrhea 25/31 (81) 9/35 (26) Vomiting 10/31 (32) 10/35 (6)</p>
<p>6. Moore 1989</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>	<p>NR</p>	<p>Non-protein Calorie Intake, day 5 1847 ± 123 2261 ± 60 p=0.01 Nitrogen intake, day 5, p=0.01 12.4±0.8 15.4±0.4 N balance, day 5, p=NS -0.3±0.1 0.1±0.8 Blood Sugars no difference between the groups Non-septic Complications 6/29 (21) 7/30 (23)</p>
<p>7. Kudsk 1992</p>	<p>Hospital 20.5 ± 19.9 (51)</p>	<p>Hospital 19.6 ± 18.8 (45)</p>	<p>2.8 ± 4.9 (51)</p>	<p>3.2 ± 6.7 (45)</p>	<p>NR</p>	<p>NR</p>	<p>NON PROTEIN Calorie Intake (kcal/kg/day) 15.7 ± 4.2 19.1 ± 3.3 p<0.05 Diarrhea 11/51 (22) 7/45 (16)</p>

8. Dunham 1994	NR	NR	NR	NR	NR	NR	<p>Calorie Intake no difference between the groups</p> <p>Protein Intake no difference between the groups</p> <p>Nutrition-related Complications 3/12 (25) 2/15 (13)</p>
9. Borzotta 1994	Hospital (assumed) 39 ± 23.1	Hospital (assumed) 36.9 ± 14	NR	NR	\$121,941	\$112,450	<p>Calorie Intake no difference between the groups</p> <p>Placement Complications 3/28 (11) 0/21 (0)</p> <p>Aspiration 3/28 (11) 0/21 (0)</p> <p>Hyperglycemia 12/28 (44) 16/21 (76)</p> <p>P=<0.05</p> <p>Diarrhea 30% 62%</p>
10. Hadfield 1995	NR	NR	NR	NR	NR	NR	NR
11. Kalfarentzos 1997	ICU 11 (5-21) * Hospital 40 (25-83) *	ICU 12 (5-24) * Hospital 39 (22-73) *	15 (6-16) *	11 (7-31) *	£70/day savings	NR	<p>Non-protein Calorie Intake (kcal/kg/day) 24.1 24.5</p> <p>p=NS</p> <p>Protein Intake (gm/kg/day) 1.43 1.45</p> <p>p=NS</p> <p>Hyperglycemia 4/18 (22) 9/20 (45)</p> <p>p=NR</p>
12. Woodcock 2001	33.2 ± 43 (16)	27.3 ± 18.7 (18)	NR	NR	NR	NR	<p>% Target Intake Achieved 54.1% 96.7%</p> <p>p<0.001</p> <p>< 80% Target Intake 62.5% 6.3%</p> <p>p<0.001</p>
13. Casas 2007	Hospital 30.2 (average)	Hospital 30.7 (average)	NR	NR	NR	NR	<p>Kcal/kg/d, p=ns, n=11 in both groups 20.09±1.83 20.8±1.68</p> <p>P=NS</p> <p>Nitrogen g/kg/d, p<0.005 0.148±0.016 0.186±0.009</p>

14. Chen 2011	ICU 9.09 ± 2.75 Hospital 23.32 ± 5.6	ICU 9.60 ± 3.06 Hospital 22.24 ± 3.27	7.95 ± 2.11	8.23 ± 2.42	NR	NR	Non-infectious Complications 10/49 (20) 21/49 (43) Gastric Residuals 6/49 (12) 0/49 (0) Diarrhea 6/49 (12) 8/49 (16)
15. Meirelles 2011	ICU 14 (5-26)	ICU 14 (6-24)	NR	NR	NR	NR	Kcal over 5 days 5958 +/- 3619 6586 +/- 1052 P=0.34 Mean daily N-balance, p=0.34 -4.6g/day -5.9g/day Blood Glucose (mg/dl) 102.4 (91.6 – 113.2) 134.4 (122.6-146.2) p < 0.0111
16. Wang 2013	NR	NR	NR	NR	NR	NR	NR
17. Sun 2013	ICU 9 (5-14)	ICU 12 (8-21)	NR	NR	NR	NR	NR
18. Harvey 2014	ICU 11.3 ± 12.5 (1197) Hospital 26.8 ± 33.2 (1186)	ICU 12 ± 13.5 (1190) Hospital 27.5 ± 33.9 (1185)	8.2 ± 9.3 (1197)	8.7 ± 11.5 (1189)	NR	NR	Vomiting 1/1197 (0.1) 1/1197 (0.1) Aspiration/Regurgitation 4/1197 (0.3) 2/1191 (0.2) Diarrhea 250/1197 (21) 192/1191 (16.2) Total kcal received during intervention period (kcal/kg) 74 ± 44 89 ± 44 P=NR Total protein received during intervention period (g/kg) 3 ± 2 3 ± 2
19. Xi 2014	ICU 8.52 ± 3.6 (22) Hospital 20.43 ± 10.49 (22)	ICU 20.33 ± 4.47 (23) Hospital 38.76 ± 15.04 (23)	2.96 ± 1.74 (22)	8.62 ± 3.6 (23)	Hospital cost x \$10⁴ 1.45 ± 0.25	Hospital cost x \$10⁴ 3.47 ± 0.69	NR

20. Reigner 2017	ICU 9.0 (5.0-16.0) 13.7±16.1** N=1201 Hospital 17.0 (8.0-32.0) 25.1±28.4** N=1202	ICU 10.0 (5.0-17.0) 13.7±13.9** N=1207 Hospital 18.0 (9.0-33.0) 25.9±27.0** N=1208	10.7±14.4** N=1201	10.9±12.6** N=1207	NR	NR	Diarrhea 432/1202 (36) 393/1208 (33) Kcal/kg/d 17.8 ± 5.5 19.6 ± 5.3 P<0.0001 Protein g/kg/d 0.7 ± 0.2 0.8 ± 0.2 P<0.0001
21) Chen 2019	Hospital 16.09 ± 1.64	Hospital 27.16 ± 4.25	NR	NR	NR	NR	Total Protein level significantly higher increases in EN vs PN p=0.032 Albumin levels significantly higher increases in EN vs PN p=0.028

C.Random: concealed randomization; ITT: intent to treat
 ± () : mean ± Standard deviation (number)
 * median/mean values, no standard deviation hence not included in meta-analysis
 NR: not reported reported data pertaining to ICU patients only
 ‡ refers to the # of patients with infections unless specified
 † presumed hospital mortality unless otherwise specified
 NS = not statistically significant
 ** data obtained directly from authors

Figure 1. Studies comparing EN vs PN: Overall Mortality




Figure 2. Overall mortality in studies with hyperglycemia where the PN group had higher blood sugars than the EN group




Figure 3. ICU Mortality




Figure 4. Studies comparing EN vs PN: Infectious complications




Figure 5. Infections in studies with hyperglycemia where the PN group had higher blood sugars than the EN group



Figure 6. Hospital LOS




Figure 7. ICU LOS





Figure 8. Mechanical Ventilation



References

Included Studies

1. Rapp RP, Young DB, Twyman D. The favorable effect of early parenteral feeding on survival in head-injured patients. *J Neurosurgery* 1983;58:906-12.
2. Adams S, Dellinger EP, Wertz MJ. Enteral versus parenteral nutritional support following laparotomy for trauma: A randomized prospective trial. *J Trauma* 1986;26(10):882-891.
3. Young B, Ott L, Twyman D et al. The effect of nutritional support on outcome from severe head injury. *J Neurosurg* 1987;67:668-76.
4. Cerra FB, McPherson JP, Konstantinides FN, Konstantinides NN, Teasley KM. Enteral nutrition does not prevent multiple organ failure syndrome (MOFS) after sepsis. *Surgery* 1988;104:727-33.
5. Peterson VM, Moore EE, Jones TN, Rundus C, Emmett M, Moore FA, McCroskey BL, Haddix T, Parsons PE. Total enteral nutrition versus total parenteral nutrition after major torso injury: attenuation of hepatic protein reprioritization. *Surgery* 1988 Aug;104(2):199-207.
6. Moore FA, Moore EE, Jones TN, McCroskey BL, Peterson VM. TEN versus TPN following major abdominal trauma - Reduced septic morbidity. *J Trauma* 1989;29:916-923.
7. Kudsk KA, Croce MA, Fabian TC et al. Enteral versus parenteral feeding: Effects on septic morbidity after blunt and penetrating abdominal trauma. *Ann Surg* 1992;215:503-13.
8. Borzotta AP, Pennings J, Papisadero B et al. Enteral versus parenteral nutrition after severe closed head injury. *J Trauma* 1994;37(3):459-468.
9. Dunham CM, Frankenfield D, Belzberg H, Wiles C, Cushing B, Grant Z. Gut failure-predictor of or contributor to mortality in mechanically ventilated blunt trauma patients? *J Trauma* 1994;37(1):30-34.
10. Hadfield RJ, Sinclair DG, Houldsworth PE, Evans TW. Effects of enteral and parenteral nutrition on gut mucosal permeability in the critically ill. *Am J Respi Crit Care Med* 1995;152:1545-8.
11. Kalfarentzos F, Kehagias J, Mead N, Kokkinis K, Gogos CA. Enteral nutrition is superior to parenteral nutrition in severe acute pancreatitis: Results of a randomized prospective trial. *British J Surg* 1997;84:1665-9.
12. Woodcock NP, Zeigler D, Palmer MD, Buckley P, Mitchell CJ, Macfie J. Enteral versus parenteral nutrition: A pragmatic study. *Nutrition* 2001;17:1-12.
13. Casas M, Mora J, Fort E, Aracil C, Busquets D, Galter S, Jáuregui CE, Ayala E, Cardona D, Gich I, Farré A. [Total enteral nutrition vs. total parenteral nutrition in patients with severe acute pancreatitis] [Article in Spanish] *Rev Esp Enferm Dig*. 2007 May;99(5):264-9.
14. Chen F, Wang J, Jiang Y. Influence of different routes of nutrition on the respiratory muscle strength and outcome of elderly patients in respiratory intensive care unit. *Chinese Journal of Clinical Nutrition*. 2011;1:7-11.
15. Justo Meirelles CM, de Aguiar-Nascimento JE. Enteral or parenteral nutrition in traumatic brain injury: a prospective randomised trial. *Nutr Hosp*. 2011 Sep-Oct;26(5):1120-4.
16. Wang G, Wen J, Xu L, Zhou S, Gong M, Wen P, Xiao X. Effect of enteral nutrition and ecoinutrition on bacterial translocation and cytokine production in patients with severe acute pancreatitis. *J Surg Res*. 2013 Aug;183(2):592-7.
17. Sun JK, Mu XW, Li WQ, Tong ZH, Li J, Zheng SY. Effects of early enteral nutrition on immune function of severe acute pancreatitis patients. *World J Gastroenterol*. 2013 Feb 14;19(6):917-22.
18. Harvey SE, Parrott F, Harrison DA, et al. Trial of the route of early nutritional support in critically ill adults. *N Engl J Med*. 2014;371(18):1673-1684. doi:10.1056/NEJMoa1409860
19. Xi F, Li N, Geng Y, Gao T, Zhang J, Jun T, Lin Z, Li W, Zhu W, Yu W, Li J. Effects of Delayed Enteral Nutrition on Inflammatory Responses and Immune Function Competence in Critically Ill Patients with Prolonged Fasting. *Hepatogastroenterology*. 2014 May;61(131):606-12.
20. Reignier J, Boisramé-Helms J, Brisard L, et al. Enteral versus parenteral early nutrition in ventilated adults with shock: a randomised, controlled, multicentre, open-label, parallel-group study (NUTRIREA-2). *Lancet*. 2018;391(10116):133-143.
21. Chen Y-J, Zhuang Y-D, Cai Z, Zhang Y-N, Guo S-R. Effects of enteral nutrition on pro-inflammatory factors and intestinal barrier function in patients with acute severe pancreatitis. *European Journal of Inflammation*. January 2019. doi:10.1177/2058739219827212

Excluded Studies	Reasons
Lim ST, Choa RG, Lam KH, Wong J, Ong GB. Total parenteral nutrition versus gastrostomy in the preoperative preparation of patients with carcinoma of the oesophagus. <i>Br J Surg</i> . 1981 Feb;68(2):69-72.	Not ICU patients
Sako K, Loré JM, Kaufman S, Razack MS, Bakamjian V, Reese P. Parenteral hyperalimentation in surgical patients with head and neck cancer: a randomized study. <i>J Surg Oncol</i> . 1981;16(4):391-402.	Not ICU patients
Bauer E, Graber R, Brodike R et al. Ernährungsphysiologische, immunologische und klinische parameter bei prospektiv randomisierten patienten unter enteraler oder parenteraler ernährungstherapie nach dickdarmoperationen. <i>Infusionstherapie</i> 1984;11:165-167.	Unclear if ICU patients
Quayle AR, Mangnall D, Clark RG. A comparison of immediate post-operative enteral and parenteral nutrition in patients with gastric carcinoma. <i>Clin Nutr</i> 1984;3:35-39.	Not critically ill patients
Seri S, Aquilio E. Effects of early nutritional support in patients with abdominal trauma. <i>It J Surg Sci</i> 1984;14:223-7.	Excluded April 2002 as not likely ICU patients
Wiedeck H, Merkle N, Herfarth Ch, Grunert A. Postoperative enteral nutrition following resection of the colon. <i>Anaesthesist</i> 1984;33:63-67.	Elective surgery patients
Costalat G, Vernhet J. Nutrition enterale postoperative precoce par catheter jejunal en chirurgie digestive lourde. Comparison avec la nutrition parenterale exclusive. <i>Chirurgie</i> 1985 ;111 :708-714.	Elective surgery patients
Bower RH, Talamini MA, Sax HC. Postoperative enteral vs parenteral nutrition: A randomized controlled trial. <i>Arch Surg</i> 1986;121:1040-5.	Elective surgery patients
Fletcher JP, Little JM. A comparison of parenteral nutrition and early postoperative enteral feeding on the nitrogen balance after major surgery. <i>Surgery</i> 1986;100:21-4.	Surgical patients
Hadley MN, Gramh TW, Harrington T. Nutritional support and neurotrauma: A critical review of early nutrition in forty-five acute head injury patients. <i>Neurosurgery</i> 1986;19:367-73.	Pseudo-randomized
Young B, Ott L, Haack D. Effect of total parenteral nutrition upon intracranial pressure in severe head injury. <i>J Neurosurg</i> 1987;67:76-80.	No clinical outcomes
Greenberg GR, Fleming CR, Jeejeebhoy KN, Rosenberg IH, Sales D, Tremaine WJ. Controlled trial of bowel rest and nutritional support in the management of Crohn's disease. <i>Gut</i> . 1988 Oct;29(10):1309-15.	Not ICU patients
Hamaoui E, Lefkowitz R, Olender L et al. Enteral nutrition in the early postoperative period: A new semi-elemental formula versus total parenteral nutrition. <i>JPEN:J Parenter Enteral Nutr</i> 1990;14:501-7.	Elective surgery patients
Moore FA, Feliciano DV, Andrassy RJ et al. Early enteral feeding, compared with parenteral, reduces postoperative septic complications: The results of a meta-analysis. <i>Ann Surg</i> 1992;216:172-83.	Meta-analysis
Von Meyenfeldt MF, Meijerink WJ, Rouflart MM, Builmaassen MT, Soeters PB.C. Perioperative nutritional support: a randomised clinical trial. <i>lin Nutr</i> . 1992 Aug;11(4):180-6.	Elective surgery patients
González-Huix F, Fernández-Bañares F, Esteve-Comas M, Abad-Lacruz A, Cabré E, Acero D, Figa M, Guilera M, Humbert P, de León R, et al. Enteral versus parenteral nutrition as adjunct therapy in acute ulcerative colitis. <i>Am J Gastroenterol</i> . 1993 Feb;88(2):227-32.	Not ICU patients
Iovinelli G, Marsili I, Varrassi G. Nutrition support after total laryngectomy. <i>JPEN J Parenter Enteral Nutr</i> . 1993 Sep-Oct;17(5):445-8.	Elective surgery patients
Kudsk KA. Gut mucosal nutritional support - Enteral nutrition as primary therapy after multiple system trauma. <i>Gut</i> 1994;35:S52-S54.	Duplicate study of '92
Wicks C, Somasundaram S, Bjarnason I et al. Comparison of enteral feeding and total parenteral nutrition after liver transplantation. <i>Lancet</i> 1994;344(8926):837-40.	Elective surgery patients
Baigrie RJ, Devitt PG, Watkin DS. Enteral versus parenteral nutrition after oesophagogastric surgery: a prospective randomized comparison. <i>Aust N Z J Surg</i> . 1996 Oct;66(10):668-70.	Elective surgery patients
Hernandez-Aranda JC, Gallo-Chico B, Ramirez-Barba EJ. Nutritional support in severe acute pancreatitis. Controlled clinical trial. <i>Nutr Hosp</i> 1996;11(3):160-6.	Not RCT

Suchner U, Senftleben U, Eckart T et al. Enteral versus parenteral nutrition: Effects on gastrointestinal function and metabolism. <i>Nutrition</i> 1996;12:13-22.	No clinical outcomes
Georgiannos SN, Renaut AJ, Goode AW. Short-term restorative nutrition in malnourished patients: Pro's and con's of intravenous and enteral alimentation using compositionally matched nutrients. <i>Int Surg</i> 1997;82:301-306.	Not ICU patients
Gianotti L, Braga M, Vignali A, Balzano G, Zerbi A, Bisagni P, Di Carlo V. Effect of route of delivery and formulation of postoperative nutritional support in patients undergoing major operations for malignant neoplasms. <i>Arch Surg.</i> 1997 Nov;132(11):1222-9.	Not ICU patients
McClave SA, Greene LM, Snider HL, Makk LJ, Cheadle WG, Owens NA, Dukes LG, Goldsmith LJ. Comparison of the safety of early enteral vs parenteral nutrition in mild acute pancreatitis. <i>JPEN J Parenter Enteral Nutr.</i> 1997 Jan-Feb;21(1):14-20.	Not ICU patients
Reynolds JV, Kanwar S, Welsh FK, Windsor AC, Murchan P, Barclay GR, Guillou PJ. Does the route of feeding modify gut barrier function and clinical outcome in patients after major upper gastrointestinal surgery? <i>JPEN J Parenter Enteral Nutr.</i> 1997 Jul-Aug;21(4):196-201.	Not ICU patients
Sand J, Luostarinen M, Matikainen M. Enteral or parenteral feeding after total gastrectomy: prospective randomised pilot study. <i>Eur J Surg.</i> 1997 Oct;163(10):761-6.	Not ICU patients
Shirabe K, Matsumata T, Shimada M, Takenaka K, Kawahara N, Yamamoto K, Nishizaki T, Sugimachi K. A comparison of parenteral hyperalimentation and early enteral feeding regarding systemic immunity after major hepatic resection--the results of a randomized prospective study. <i>Hepatogastroenterology.</i> 1997 Jan-Feb;44(13):205-9.	Not ICU patients
Braga M, Gianotti L, Vignali A, Cestari A, Bisagni P, Di C, V. Artificial nutrition after major abdominal surgery: Impact of route of administration and composition of the diet. <i>Crit Care Med</i> 1998;26:24-30.	Elective surgery patients
Windsor ACJ, Kanwar S, Li AGK et al. Compared with parenteral nutrition, enteral feeding attenuates the acute phase response and improves disease severity in acute pancreatitis. <i>Gut</i> 1998;42:431-5.	Not ICU patients
Oláh A, Pardavi G, Belágyi T. [Early jejunal feeding in acute pancreatitis: prevention of septic complications and multiorgan failure][Article in Hungarian] <i>Magy Seb.</i> 2000 Feb;53(1):7-12.	Elective surgery patients
Bozzetti F, Braga M, Gianotti L, Gavazzi C, Mariani L. Postoperative enteral versus parenteral nutrition in malnourished patients with gastrointestinal cancer: A randomised multicentre trial. <i>Lancet</i> 2001;358:1487-92.	Elective surgery patients
Braga M, Gianotti L, Gentilini O, Parisi V, Salis C, Di C, V. Early postoperative enteral nutrition improves gut oxygenation and reduces costs compared with total parenteral nutrition. <i>Crit Care Med</i> 2001;29:242-8.	Elective surgery patients
Braunschweig CL, Levy P, Sheean PM, Wang X. Enteral compared with parenteral nutrition: A meta-analysis. <i>Am J Clin Nutr</i> 2001;74:534-42.	Meta-analysis
Heyland DK, Montalvo M, MacDonald S et al. Total parenteral nutrition in the surgical patient: a meta-analysis. <i>Can J Surg</i> 2001;44(2):102-111.	Meta-analysis
Pacelli F, Bossola M, Papa V et al. Enteral vs parenteral nutrition after major abdominal surgery: An even match. <i>Arch Surg</i> 2001;136:933-6.	Elective surgery patients
Abou-Assi S, Craig K, O'Keefe SJ. Hypocaloric jejunal feeding is better than total parenteral nutrition in acute pancreatitis : results of a randomized comparative study. <i>Am J Gastroenterology</i> 2002;97(9):2255-2262.	Not ICU patients
Huang YC, Yen CE, Cheng CH, Jih KS, Kan MN. Nutritional status of mechanically ventilated critically ill patients: comparison of different types of nutritional support. <i>Clin Nutr</i> 2002;101-7.	Not RCT
Olah A, Pardavi G, Belágyi T, Nagy A, Issekutz A, Mohamed GE. Early nasojejunal feeding in acute pancreatitis is associated with a lower complication rate. <i>Nutrition</i> 2002;18(3):259-62.	Pseudo-randomized
Rayes N, Hansen S, Seehofer D, Müller AR, Serke S, Bengmark S, Neuhaus P. Early enteral supply of fiber and Lactobacilli versus conventional nutrition: a controlled trial in patients with major abdominal surgery. <i>Nutrition.</i> 2002 Jul-Aug;18(7-8):609-15.	Elective surgery patients

Gupta R, Patel K, Calder PC, Yaqoob P, Primrose JN, Johnson CD. A randomised clinical trial to assess the effect of total enteral and total parenteral nutritional support on metabolic, inflammatory and oxidative markers in patients with predicted severe acute pancreatitis (APACHE II > or =6). <i>Pancreatolgy</i> . 2003;3(5):406-13.	Not ICU patients
Zhao G, Wang CY, Wang F, Xiong JX. Clinical study on nutrition support in patients with severe acute pancreatitis. <i>World journal of gastroenterology: WJG</i> 2003;9(9):2105-8.	No clinical outcomes
Louie BE, Noseworthy T, Hailey D, Gramlich LM, Jacobs P, Warnock GL. 2004 MacLean-Mueller prize enteral or parenteral nutrition for severe pancreatitis: a randomized controlled trial and health technology assessment. <i>Can J Surg</i> . 2005 Aug;48(4):298-306.	Not ICU patients
Peter JV, Moran JL, Phillips-Hughes J. A meta-analysis of treatment outcomes of early enteral versus early parenteral nutrition in hospitalized patients. <i>Crit Care Med</i> . 2005 Jan;33(1):213-20.	Meta-analysis
Simpson F, Doig GS. Parenteral vs. enteral nutrition in the critically ill patient: a meta-analysis of trials using the intention to treat principle. <i>Intensive Care Med</i> . 2005 Jan;31(1):12-23.	Meta-analysis
Eckerwall GE, Axelsson JB, Andersson RG. Early nasogastric feeding in predicted severe acute pancreatitis. A clinical, randomized study. <i>Ann Surg</i> 2006;244(6):959-967.	Not ICU patients
Petrov MS, Kukosh MV, Emelyanov NV. A randomized controlled trial of enteral versus parenteral feeding in patients with predicted severe acute pancreatitis shows a significant reduction in mortality and in infected pancreatic complications with total enteral nutrition. <i>Dig Surg</i> . 2006;23(5-6):336-44; discussion 344-5.	Unable to confirm if patients were in ICU. Contacted authors but did not get needed details
Radrizzani D, Bertolini G, Facchini R, Simini B, Bruzzone P, Zanforlin G, et al. Early enteral immunonutrition vs. parenteral nutrition in critically ill patients without severe sepsis: a randomized clinical trial. <i>Intensive care medicine</i> 2006;32(8):1191-8.	Control group received nonstandard EN formula (arginine)
Chen Z, Wang S, Yu B, Li A. A comparison study between early enteral nutrition and parenteral nutrition in severe burn patients. <i>Burns</i> 2007;33(6):708-12.	No clinical outcomes
Petrov MS, Zagainov VE. Influence of enteral versus parenteral nutrition on blood glucose control in acute pancreatitis: a systematic review. <i>Clin Nutr</i> . 2007 Oct;26(5):514-23.	Systematic review
Tian. [The morphological alterations of jejunal mucosa accepting early enteral nutrition for post-operative patients with severe acute pancreatitis]. <i>Sichuan da xue xue bao Yi xue ban = Journal of Sichuan University Medical science edition</i> 2007;38(2):264-7. https://www.researchgate.net/publication/298861455_Enteral_versus_parenteral_nutrition_in_mechanically_ventilated_patients	Abstract, unable to download, included in Cochrane review above but they only included abstract data
Cao Y, Xu Y, Lu T, Gao F, Mo Z. Meta-analysis of enteral nutrition versus total parenteral nutrition in patients with severe acute pancreatitis. <i>Annals of Nutrition and Metabolism</i> 2008;53(3-4):268-75.	Meta-analysis
Farimani M, Bajestani N. Comparison of early enteral feeding versus parenteral nutrition after resection of esophageal cancer. <i>Journal of Critical Care</i> . 2008; 23(3):448	Elective surgery patients
Lam NN, Tien NG, Khoa CM. Early enteral feeding for burned patients--an effective method which should be encouraged in developing countries. <i>Burns</i> . 2008 Mar;34(2):192-6.	Pseudo-randomized
Petrov MS, Pylypchuk RD, Emelyanov NV. Systematic review: nutritional support in acute pancreatitis. <i>Aliment Pharmacol Ther</i> . 2008 Sep 15;28(6):704-12.	Systematic review

Cheng XT, Li SL, Liu GL, Yang XM, Lu J. [Effect of nutritional support on immune function in patients with severe pulmonary infection after renal transplantation]. <i>Nan fang yi ke da xue xue bao = Journal of Southern Medical University</i> 2009;29(6):1159-62.	Not ICU patients
Doley RP, Yadav TD, Wig JD, Kochhar R, Singh G, Bharathy KG, Kudari A, Gupta R, Gupta V, Poornachandra KS, Dutta U, Vaishnavi C. Enteral nutrition in severe acute pancreatitis. <i>JOP</i> . 2009 Mar 9;10(2):157-62.	Pseudo-randomized
Nagata S, Fukuzawa K, Iwashita Y, Kabashima A, Kinoshita T, Wakasugi K, Maehara Y. Comparison of enteral nutrition with combined enteral and parenteral nutrition in post-pancreaticoduodenectomy patients: a pilot study. <i>Nutr J</i> . 2009 Jun 11;8:24.	Elective surgery patients
Ryu J, Nam BH, Jung YS. Clinical outcomes comparing parenteral and nasogastric tube nutrition after laryngeal and pharyngeal cancer surgery. <i>Dysphagia</i> . 2009 Dec;24(4):378-86.	Elective surgery patients
Vieira JP, Araujo GF, Azevedo JR, Goldenberg A, Linhares MM. Parenteral nutrition versus enteral nutrition in severe acute pancreatitis. <i>Acta cirurgica brasileira / Sociedade Brasileira para Desenvolvimento Pesquisa em Cirurgia</i> 2010;25(5):449-54.	Not RCT
Gencer A, Ozdemir Y, Sucullu I, Filiz AI, Yucel E, Akin ML, et al. The effects of enteral immunonutrient products and total parenteral nutrition in patients who underwent major abdominal surgery [Majör abdominal kanser cerrahisi uygulanan hastalarda total parenteral nutrisyon ve enteral immunonutrisyon kar ila t r lmas]. <i>Trakya Universitesi Tip Fakultesi Dergisi</i> 2010;27(4):404-10. DOI: 10.5174/tutfd.2009.02426.1	Elective surgery patients
Altintas ND, Aydin K, Turkoglu MA, Abbasoglu O, Topeli A. Effect of enteral versus parenteral nutrition on outcome of medical patients requiring mechanical ventilation. <i>Nutrition in clinical practice : official publication of the American Society for Parenteral and Enteral Nutrition</i> 2011;26(3):322-9.	Pseudo-randomized
Cangelosi MJ, Auerbach HR, Cohen JT. A clinical and economic evaluation of enteral nutrition. <i>Current medical research and opinion</i> 2011;27(2):413-22.	Meta-analysis
Klek S, Sierzega M, Turczynowski L, szykinski P, Sczcepanek K, Kulig J. (2011). Enteral and Parenteral Nutrition in the Conservative Treatment of Pancreatic Fistula: A Randomized Clinical Trial. 2011;141:157-163.	Elective surgery patients
Wang X, Dong Y, Han X, Qi X-Q, Huang C-G, Hou L. (2013) Nutritional Support for Patients Sustaining Traumatic Brain Injury: A Systematic Review and Meta-Analysis of Prospective Studies. <i>PLoS ONE</i> . 8(3): e58838.	Meta-analysis
Boelens PG, Heesakkers FF, Luyer MD, et al. Reduction of postoperative ileus by early enteral nutrition in patients undergoing major rectal surgery: prospective, randomized, controlled trial. <i>Ann Surg</i> . 2014;259(4):649-655.	Not critically ill patients
Kilner T, Bidgood E, Benham-Mirando S, Krol R, Brealey D. Nutritional support and mortality in critically ill adults - A subset analysis of the calories trial. <i>Intensive Care Medicine Experimental</i> . 2015;3.	subset of CALORIES trial. No new relevant clinical outcomes. Abstract
van Barneveld KW, Smeets BJ, Heesakkers FF, Bosmans JW, Luyer MD, Wasowicz D, Bakker JA, Roos AN, Rutten HJ, Bouvy ND, Boelens PG. Beneficial Effects of Early Enteral Nutrition After Major Rectal Surgery: A Possible Role for Conditionally Essential Amino Acids? Results of a Randomized Clinical Trial. <i>Crit Care Med</i> . 2016 Jun;44(6):e353-61.	Elective surgery patients
Perinel J, Mariette C, Dousset B, Sielezneff I, Gainant A, Mabrut JY, Bin-Dorel S, Bechwaty ME, Delaunay D, Bernard L, Sauvanet A, Pocard M, Buc E, Adham M. Early Enteral Versus Total Parenteral Nutrition in Patients Undergoing Pancreaticoduodenectomy: A Randomized Multicenter Controlled Trial (Nutri-DPC). <i>Ann Surg</i> . 2016 Nov;264(5):731-737.	Not critically ill
Harvey SE, Parrott F, Harrison DA, et al. A multicentre, randomised controlled trial comparing the clinical effectiveness and cost-effectiveness of early nutritional support via the parenteral versus the enteral route in critically ill patients (CALORIES). <i>Health Technol Assess</i> . 2016;20(28):1-144.	Substudy of Harvey 2014 - adding QOL data from tables 32 and 33

Gundogan K, Dogan E, Coskun R, Muhtaroglu S, Sungur M, Ziegler T, Guven M. Association between the route of nutrition and adipokine hormones levels in critically ill patients: A pilot study. Intensive Care Medicine Experimental. Conference: 29th Annual Congress of the European Society of Intensive Care Medicine, ESICM 2016. Italy. 4 (no pagination), 2016.	no clinically significant outcomes; abstract; unclear if RCT
Fan MC, Wang QL, Fang W, Jiang Y, Li L, Sun P, et al. Early enteral combined with parenteral nutrition treatment for severe traumatic brain injury: effects on immune function, nutritional status and outcomes. Chinese Medical Science Journal 2016;31(4):213–20.	Pseudo-randomized
Lewis SR, Schofield-Robinson OJ, Alderson P, Smith AF. Enteral versus parenteral nutrition and enteral versus a combination of enteral and parenteral nutrition for adults in the intensive care unit. Cochrane Database Syst Rev. 2018 Jun 8;6:CD012276. doi: 10.1002/14651858.CD012276.pub2. Review. PubMed PMID: 29883514.	Meta-analysis
Shi J, Wei L, Huang R, Liao L. Effect of combined parenteral and enteral nutrition versus enteral nutrition alone for critically ill patients: A systematic review and meta-analysis. Medicine (Baltimore). 2018;97(41):e11874. doi:10.1097/MD.00000000000011874	Meta-analysis
Casaer MP, Mesotten D, Hermans G, et al. Early versus late parenteral nutrition in critically ill adults. N Engl J Med. 2011;365(6):506-517.	PN timing
Chiarelli AG, Ferrarello S, Piccioli A, et al. La nutrizione enterale totale verso la nutrizione mista enterale e parenterale in pazienti ricoverati in una terapia intensiva polivalente [Total enteral nutrition versus mixed enteral and parenteral nutrition in patients at an intensive care unit]. Minerva Anesthesiol. 1996;62(1-2):1-7.	Language barrier