

5.5 Strategies to Optimize the Delivery of EN: Use of and Threshold for Gastric Residual Volumes

March 2013

NEW SECTION in 2013

Recommendation: There are insufficient data to make a recommendation for not checking gastric residual volumes or a specific gastric residual volume threshold. Based on 2 level 2 studies, a gastric residual volume of either 250 or 500 mLs (or somewhere in between) is acceptable as a strategy to optimize delivery of enteral nutrition in critically ill patients.

Discussion:

The committee noted that in the multicenter study (Reignier 2013), not checking gastric residual volumes was associated with increased rates of vomiting, despite no differences in clinical outcomes. Nutritional adequacy was greater in the 'not checking GRV' group but differences were minimal (111 calories over the first week). Given the concerns about the external validity of the trial (under-represents difficult to feed patients i.e. multiorgan failure and surgical) and the signals associating vomiting from gastrointestinal intolerance with increased infection, length of stay and mortality in critically ill patients (1), the committee agreed not to make a recommendation for abandoning the practice of checking GRVs.

The committee noted that in Spanish multicentre trial (Montejo 2010), there was an absence of any clinical effect of increasing the gastric residual volume threshold and that the increase in nutritional adequacy with the higher GRV was minimal but statistically significant (84 vs. 88 % of goal calories). Despite the potential safety of an approach that used a high GRV threshold, as evidenced by the lack of increased gastrointestinal complications, concerns regarding potential microaspiration were raised. Opposing views about the risk of higher gastric residual volumes exist (2, 3). Furthermore, in the Montejo et al study, patients were predominately medical patients and there was a lack of information about their hemodynamic stability. Thus, the generalizability of the results to all ICU patients that might receive a feeding protocol in a given ICU is not clear. The committee agreed that a strong recommendation could not be made for higher GRVs of 500 mLs but it was agreed that a range of 250-500 mLs be recommended.

(1) Metheny NA, Schallom L, Oliver DA, Clouse RE. Gastric residual volume and aspiration in critically ill patients receiving gastric feedings. *Am J Crit Care* 2008;17:512-520.

(2) Mentec H, Dupont H, Bocchetti M, Cani P, Ponche F, Bleichner G. Upper digestive intolerance during enteral nutrition in critically ill patients: frequency, risk factors, and complications. *Crit Care Med*. 2001;29(10):1955-1961.

(3) McClave SA, Lukan JK, Stefater JA, et al. Poor validity of residual volumes as a marker for risk of aspiration in critically ill patients. *Crit Care Med* 2005;33(2):324-330.

Semi Quantitative Scoring

Values	Definition	2013 Score (0,1,2,3) High vs Lower	2013 Score (0,1,2,3) 250 mls vs none
Effect size	Magnitude of the absolute risk reduction attributable to the intervention listed--a higher score indicates a larger effect size	0	0
Confidence interval	95% confidence interval around the point estimate of the absolute risk reduction, or the pooled estimate (if more than one trial)--a higher score indicates a smaller confidence interval	0	0
Validity	Refers to internal validity of the study (or studies) as measured by the presence of concealed randomization, blinded outcome adjudication, an intention to treat analysis, and an explicit definition of outcomes--a higher score indicates presence of more of these features in the trials appraised	1	3
Homogeneity or Reproducibility	Similar direction of findings among trials--a higher score indicates greater similarity of direction of findings among trials	n/a	n/a
Adequacy of control group	Extent to which the control group presented standard of care (large dissimilarities=1, minor dissimilarities=2, usual care=3)	3	3
Biological Plausibility	Consistent with understanding of mechanistic and previous clinical work (large inconsistencies=1, minimal consistencies=2, very consistent=3)	2	2
Generalizability	Likelihood of trial findings being replicated in other settings (low likelihood i.e. single centre=1, moderate likelihood i.e. multicentre with limited patient population or practice setting=2, high likelihood i.e. multicentre, heterogenous patients, diverse practice settings=3)	1	1
Low cost	Estimated cost of implementing the intervention listed--a higher score indicates a lower cost to implement the intervention in an average ICU	3	3
Feasible	Ease of implementing the intervention listed--a higher score indicates greater ease of implementing the intervention in an average ICU	3	3
Safety	Estimated probability of avoiding any significant harm that may be associated with the intervention listed--a higher score indicates a lower probability of harm	2	1

5.5 Strategies to Optimize the Delivery of EN: Threshold of Gastric Residual Volumes

March 2013

Questions:

1. Does the use of higher gastric residual volume threshold (GRVs) result in better outcomes in the critically ill adult patient?
2. Does not checking gastric residual volumes compared to a GRV of 250 mL result in better outcomes in the critically ill adult patient?

Summary of evidence: There was one level 2 multicentre trial that compared a gastric residual volume of 500 mLs to 250 mLs (Montejo 2010). One study compared higher gastric residual volume threshold to lower within the context of a feeding protocol that also included motility agents (Pinilla 2001) and was included in the section 5.1 Feeding Protocols. The study by Taylor et al 1999 compared full rate EN with higher gastric residual volume thresholds vs gradual start EN with lower gastric residual volume thresholds was included in the section 3.2 Target Dose EN. There was also a multicenter trial that compared not measuring gastric residual volumes to 250 mLs (Reignier 2013).

Mortality: In the study by Montejo (2010) there were no significant difference between the two groups in ICU mortality (RR 1.25, 95% CI 0.78, 2.01, $p=0.35$) or hospital mortality (RR 1.01, 95% CI 0.74, 1.38, $p=0.94$). There were no differences in 28 day or 90 day mortality between the group that did not check gastric residual volumes vs. the group that checked GRVs > 250 ml in the multicentre study (Reignier 2013).

Infections: In the study by Montejo (2010), no significant differences were found in pneumonia between the two groups (RR 1.03, 95% CI 0.72, 1.46, $p=0.88$). There were no significant differences in ICU acquired infections or ventilator associated pneumonia rates between the group that did not check gastric residual volumes vs. the group that did check GRVs in the multicentre study (Reignier 2013).

LOS & ventilator days: In the study by Montejo (2010), there were no differences in ICU length of stay between the groups (WMD 0.90, 95% CI -2.60, 4.40, $p=0.61$) and no significant difference in duration of ventilation (WMD 0.90, 95% CI -2.02, 3.82, $p=0.55$). There were no differences in ICU or hospital length of stay between the group that did not check gastric residual volumes vs. the group that checked GRVs > 250 ml in the multicentre study (Reignier 2013).

Other: In the study by Montejo (2010), the frequency of gastrointestinal complications was significantly lower in the 500mL GRV vs 250 mLs GRV group and this was mainly due to the lower incidence of high GRVs when compared to the lower GRV group. There were no differences between these groups in the number of patients with abdominal distention ($p=0.83$), diarrhea ($p=0.95$), emesis ($p=0.31$), regurgitation ($p=0.41$) or aspiration ($p=0.48$). However, the amount of nutrition delivered in week 1 was significantly higher in the group with the 500ml GRVs threshold ($p=0.0002$). In the Reignier study, caloric target was achieved in a higher proportion of patients in the group not checking GRVs compared to the groups that did ($p<0.001$) and there was a lower cumulative calorie deficit from Day 0-7 than this group. There were higher rates of vomiting in the group that did not check gastric residual volumes but no differences in diarrhea.

Conclusions:

1. GRVs of 500 mLs vs 250 mLs have no effect on mortality, infections or ICU LOS
2. Not checking GRVs vs checking GRVs > 250 ml threshold has no effect on mortality, infections, ICU/hospital length of stay
3. GRVs of 500 mLs vs 250 mLs are not associated with increased gastrointestinal complications
4. GRVs of 500 mLs vs 250 mLs are associated with significantly better nutrition delivery.
5. Not checking GRVs vs checking GRVs > 250 ml threshold is associated with a significant better caloric delivery.

Level 1 study: if all of the following are fulfilled: concealed randomization, blinded outcome adjudication and an intention to treat analysis.

Level 2 study: If any one of the above characteristics are unfulfilled.

*p-value calculated from RevMan and differs slightly from that reported in the article.

Table 1. Randomized studies evaluating gastric residual volume in critically ill patients (continued)

Study	Length of Stay		Mechanical Ventilation		Other	
1) Montejo 2010	GRV 500mL ICU 20.7 ± 16.2 (157)	GRV 200mL ICU 19.8 ± 15.8 (165)	GRV 500mL 15.6 ± 13.6 (157)	GRV 200mL 14.7 ± 13.1 (165)	GRV 500ml GI Complications 75/157 (48)	GRV 200mL 105/165 (64)
	WMD 0.90, 95% CI -2.60, 4.40, p=0.61		WMD 0.90, 95% CI -2.02, 3.82, p=0.55		p=0.004	
					42/157 (27)	High GRV 70/165 (42) p=0.003
					16/157 (10)	Abdominal distention 18/165 (11) p=0.83
					31/157 (20)	Diarrhea 33/165 (20) p=0.95
					17/157 (11)	Emesis 24/165 (15) p=0.31
					8/157 (5)	Regurgitation 12/165 (7) p=0.41
					1.157 (1)	Aspiration 0/165 (0) p=0.48
					88.2%	Mean Diet Volume Ratio in 1 st week of EN 84.48% p=0.0002

2) Reignier 2013	No GRV	GRV 250mL	No GRV	GRV 250mL	No GRV	GRV 250mL
	10 (6-17)	10 (7-17)	7 (4-13)	7 (5-13)	90/227 (40)	60/222 (27)
	ICU				Vomiting	
	17 (9-31)	19 (10-32)			51/227 (23)	51/222 (23)
	Hospital				Diarrhea	
					90/227 (40)	141/222 (64)
					EN intolerance	

C.Random: concealed randomization
 † presumed hospital mortality unless otherwise specified

ITT: intent to treat; NA: not available
 ± () : mean ± Standard deviation (number)

‡ refers to the # of patients with infections unless specified